TI Year 4 – 6 Metrics with Methodology

Measurement Stewards Key			
CMS Core Set PEDS Adult	<u>CMS</u> <u>ScoreCard</u>	<u>Statewide</u> (STCs)	<u>NCQA</u> <u>HEDIS</u>
			•

Pediatric PCP			
Chosen Performance Measure	Measure Description		
Well child visits in third, fourth, fifth and sixth years of life	Percentage of children ages 3 to 6 who had one or more well-child visits with a primary care practitioner (PCP) during the measurement year.	••••	
Adolescent well-care visits	Percentage of adolescents ages 12 to 21 who had at least one comprehensive well-care visit with a primary care practitioner (PCP) or an obstetric/gynecologic (OB/GYN) practitioner during the measurement year.	•••	
Well-child visits in the first 15 months of life	Percentage of children who turned 15 months old during the measurement year and who had the following number of well-child visits with a primary care practitioner (PCP) during their first 15 months of life: No well-child visits Four well-child visits One well-child visits Five well-child visits Two well-child visits Six or more well-child visits	•••	

Pediatric BH		
Chosen Performance Measure	Measure Description	Measurement Stewards
Pediatric follow-up after hosp. for mental illness ages 6-17 (30 - Day)	Percentage of discharges for children ages 6 to 17 who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the child received follow-up within 30 days after discharge	•••
Pediatric follow-up after hosp. for mental illness ages 6-17 (7 - Day)	Percentage of discharges for children ages 6 to 17 who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the child received follow-up within 7 days after discharge	•••
Metabolic monitoring for children and adolescents on antipsychotics	Assesses the percentage of children and adolescents with ongoing antipsychotic medication use who had metabolic testing during the year.	•

Adult PCP & BH		
Chosen Performance Measure	Measure Description	Measurement Stewards
Follow up after hospitalization for Mental Illness: 18 and older (30 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 30 days after discharge	••••
Follow up after hospitalization for Mental Illness: 18 and older (7 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 7 days after discharge	••••
Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using antipsychotic medications	Percentage of beneficiaries ages 18 to 64 with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.	••

Justice		
Chosen Performance Measure	Measure Description	Measurement Stewards
Engagement of Alcohol and Other Drug Abuse or Dependence Treatment (34 - Day)	Percentage of beneficiaries age 18 and older with a new episode of alcohol or other drug (AOD) abuse or dependence who received continued engagement of AOD Treatment. Percentage of beneficiaries who initiated treatment and who had two or more additional AOD services or medication treatment within 34 days of the initiation visit	•••
Follow up after hospitalization for Mental Illness: 18 and older (30 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 30 days after discharge	••••
Follow up after hospitalization for Mental Illness: 18 and older (7 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 7 days after discharge	••••
Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are using antipsychotic medications	Percentage of beneficiaries ages 18 to 64 with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the measurement year.	••

Hospital		
Chosen Performance Measure	Measure Description	Measurement Stewards
Follow up after hospitalization for Mental Illness: 18 and older (30 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 30 days after discharge	••••
Follow up after hospitalization for Mental Illness: 18 and older (7 - Day)	Percentage of discharges for beneficiaries age 18 and older who were hospitalized for treatment of selected mental illness diagnoses or intentional self-harm and who had a follow-up visit with a mental health practitioner. Percentage of discharges for which the beneficiary received follow-up within 7 days after discharge	•••