

VACCINE	Birth	1 Month	2 months	4 months	6 months	8 months	9 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years	13-15 years	16 years	17-18 years
Respiratory syncytial virus (RSV-mAb [nirsevimab, clesrovimab])	1 dose depending on maternal RSV vaccination status (<8 months)					1 dose (8–19 months)												
Hepatitis B (HepB)	1 st dose	2 nd dose			3 rd dose													
Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series)			1 st dose	2 nd dose	3 rd dose*													
Diphtheria, tetanus, and acellular pertussis (DTaP <7 yrs)			1 st dose	2 nd dose	3 rd dose			4 th dose					5 th dose					
Haemophilus influenzae type b (Hib)			1 st dose	2 nd dose	3 rd dose		4 th dose											
Pneumococcal conjugate (PCV15, PCV20)			1 st dose	2 nd dose	3 rd dose		4 th dose											
Inactivated poliovirus (IPV)			1 st dose	2 nd dose	3 rd dose							4 th dose						
COVID-19 (1vCOV-mRNA, 1vCOV-aPS)					1 or 2 doses**							Yearly based on medical necessity						
Influenza					1 or 2 doses annually***									1 dose annually (9 years and above)				
Measles, mumps, and rubella (MMR)								1 st dose					2 nd dose					

VACCINE	Birth	1 Month	2 months	4 months	6 months	8 months	9 months	12 months	15 months	18 months	19-23 months	2-3 years	4-6 years	7-10 years	11-12 years	13-15 years	16 years	17-18 years
Varicella (VAR)								1 st dose					2 nd dose					
Hepatitis A (HepA)								2 dose series										
Tetanus, diphtheria, and acellular pertussis (Tdap ≥7 yrs)															1 dose			
Human papillomavirus (HPV)															2 dose series			
Meningococcal ACWY															1 st dose		2 nd dose	
Meningococcal B																	1 dose	

This routine pediatric vaccine schedule is reflective of evidence-based guidelines around the recommended standard timing of child and adolescent vaccines. For further information on vaccine interval recommendations, as well as additional tables listing timings for catch-up vaccines and guidelines around specific medical indications and contraindications, please refer to the most current immunization tables which can be found at: <https://downloads.aap.org/AAP/PDF/AAP-Immunization-Schedule.pdf>

For adult vaccine coverage, refer to AMPM Policy 310-M.

- * Rotavirus vaccination is a 2 or 3 dose series depending on the vaccine manufacturer. If any dose in the series is of unknown type, default to 3-dose series.
- ** 1-2 doses depending on COVID-19 vaccine manufacturer guidelines. See American Academy of Pediatrics (AAP) vaccine schedule for further guidance.
- *** For ages 6 months to 8 years 2 doses of influenza are needed for the first year of vaccination only.